

In the Claims:

39. (Currently amended) An isolated polypeptide having at least 80% amino acid sequence identity to:

- (a) the amino acid sequence of the polypeptide shown in Figure 102 (SEQ ID NO: 290);
- (b) the amino acid sequence of the polypeptide shown in Figure 102 (SEQ ID NO: 290), lacking its associated signal peptide;
- (c) the amino acid sequence of the extracellular domain of the polypeptide shown in Figure 102 (SEQ ID NO: 290); or
- (d) the amino acid sequence of the polypeptide encoded by the full-length coding sequence of the cDNA deposited under ATCC accession number 209927,  
wherein said polypeptide induces proliferation of stimulated T lymphocytes in a mixed lymphocyte reaction.

40. (Currently amended) An isolated polypeptide having at least 85% amino acid sequence identity to:

- (a) the amino acid sequence of the polypeptide shown in Figure 102 (SEQ ID NO: 290);
- (b) the amino acid sequence of the polypeptide shown in Figure 102 (SEQ ID NO: 290), lacking its associated signal peptide;
- (c) the amino acid sequence of the extracellular domain of the polypeptide shown in Figure 102 (SEQ ID NO: 290); or
- (d) the amino acid sequence of the polypeptide encoded by the full-length coding sequence of the cDNA deposited under ATCC accession number 209927,  
wherein said polypeptide induces proliferation of stimulated T lymphocytes in a mixed lymphocyte reaction.

41. (Currently amended) An isolated polypeptide having at least 90% amino acid sequence identity to:

- (a) the amino acid sequence of the polypeptide shown in Figure 102 (SEQ ID NO: 290);
- (b) the amino acid sequence of the polypeptide shown in Figure 102 (SEQ ID NO: 290), lacking its associated signal peptide;
- (c) the amino acid sequence of the extracellular domain of the polypeptide shown in Figure 102 (SEQ ID NO: 290); or
- (d) the amino acid sequence of the polypeptide encoded by the full-length coding sequence of the cDNA deposited under ATCC accession number 209927,  
wherein said polypeptide induces proliferation of stimulated T lymphocytes in a mixed lymphocyte reaction.

42. (Currently amended) An isolated polypeptide having at least 95% amino acid sequence identity to:

- (a) the amino acid sequence of the polypeptide shown in Figure 102 (SEQ ID NO: 290);
- (b) the amino acid sequence of the polypeptide shown in Figure 102 (SEQ ID NO: 290), lacking its associated signal peptide;
- (c) the amino acid sequence of the extracellular domain of the polypeptide shown in Figure 102 (SEQ ID NO: 290); or
- (d) the amino acid sequence of the polypeptide encoded by the full-length coding sequence of the cDNA deposited under ATCC accession number 209927,  
wherein said polypeptide induces proliferation of stimulated T lymphocytes in a mixed lymphocyte reaction.

43. (Currently amended) An isolated polypeptide having at least 99% amino acid sequence identity to:

- (a) the amino acid sequence of the polypeptide shown in Figure 102 (SEQ ID NO: 290);

- (b) the amino acid sequence of the polypeptide shown in Figure 102 (SEQ ID NO: 290), lacking its associated signal peptide;
- (c) the amino acid sequence of the extracellular domain of the polypeptide shown in Figure 102 (SEQ ID NO: 290); or
- (d) the amino acid sequence of the polypeptide encoded by the full-length coding sequence of the cDNA deposited under ATCC accession number 209927,  
wherein said polypeptide induces proliferation of stimulated T lymphocytes in a mixed lymphocyte reaction.

44. (Currently amended) An isolated polypeptide comprising:

- (a) the amino acid sequence of the polypeptide shown in Figure 102 (SEQ ID NO: 290);
- (b) the amino acid sequence of the polypeptide shown in Figure 102 (SEQ ID NO: 290), lacking its associated signal peptide;
- (c) the amino acid sequence of the extracellular domain of the polypeptide shown in Figure 102 (SEQ ID NO: 290); or
- (d) the amino acid sequence of the polypeptide encoded by the full-length coding sequence of the cDNA deposited under ATCC accession number 209927,  
wherein said polypeptide induces proliferation of stimulated T lymphocytes in a mixed lymphocyte reaction.

45. (Previously added) The isolated polypeptide of Claim 44 comprising the amino acid sequence of the polypeptide shown in Figure 102 (SEQ ID NO: 290).

46. (Previously added) The isolated polypeptide of Claim 44 comprising the amino acid sequence of the polypeptide shown in Figure 102 (SEQ ID NO: 290), lacking its associated signal peptide.

47. (Previously added) The isolated polypeptide of Claim 44 comprising the amino acid sequence of the extracellular domain of the polypeptide shown in Figure 102 (SEQ ID NO: 290).

48. (Previously canceled).

49. (Previously added) The isolated polypeptide of Claim 44 comprising the amino acid sequence of the polypeptide encoded by the full-length coding sequence of the cDNA deposited under ATCC accession number 209927.

50. (Previously added) A chimeric polypeptide comprising a polypeptide according to Claim 39 fused to a heterologous polypeptide.

51. (Previously added) The chimeric polypeptide of Claim 50, wherein said heterologous polypeptide is an epitope tag or an Fc region of an immunoglobulin.